

Original article

## Effects of neuromuscular electrical stimulation on the range of motion recovery in hand proximal interphalangeal sprain

## Effets de l'électrostimulation neuromusculaire sur la récupération des amplitudes articulaires dans les entorses de l'articulation interphalangienne proximale de la main

M. Boisgontier<sup>a,\*,b,c,d</sup>, N. Vuillerme<sup>b</sup>, D. Thomas<sup>c</sup>, N. Pinsault<sup>a,d</sup>,  
M. Emprin<sup>c</sup>, J.-L. Caillat-Miousse<sup>d</sup>

<sup>a</sup> Laboratoire TIMC-IMAG, UMR-UJF-CNRS 5525, faculté de médecine, bâtiment Jean-Roget, 38706 La Tronche cedex, France

<sup>b</sup> MGH Institute of Health Professions, Charlestown-Navy-Yard, 36, 1st Avenue, Boston, MA 02129-4557, United States

<sup>c</sup> Centre grenoblois de rééducation de la main et du membre supérieur, 1, boulevard Clémenceau, 38100 Grenoble, France

<sup>d</sup> École de kinésithérapie, CHU de Grenoble, BP 217, 38049 Grenoble cedex 09, France

Received 5 April 2008; accepted 10 September 2008

Available online 26 October 2008

---

### Abstract

**Aim.** – To compare the effects of active range of motion versus neuromuscular electrical stimulation superimposed to active range of motion on the range of motion recovery at the proximal interphalangeal joint following sprain.

**Patients and methods.** – Twenty patients in need of physical therapy to recover proximal interphalangeal range of motion participated voluntarily. Ranges of motion at the proximal interphalangeal joint were measured before and after each treatment with a finger goniometer.

**Results.** – Both treatments allow recuperation of the range of motion on the proximal interphalangeal joint. Moreover, the effectiveness of the superimposed technique in improving the range of motion is significantly better than that observed after active range of motion on its own.

**Conclusion.** – These findings highlighted the superimposed technique as an effective method, which could be integrated in rehabilitation protocols for recovering the proximal interphalangeal joint range of motion following sprain.

© 2008 Elsevier Masson SAS. All rights reserved.

### Résumé

**Objectif.** – Comparer les effets de la mobilisation active volontaire à ceux de l'électrostimulation neuromusculaire surimposée à la mobilisation active volontaire sur la récupération des amplitudes de l'articulation interphalangienne proximale suite à une entorse.

**Patients et méthodes.** – Vingt sujets nécessitant une prise en charge kinésithérapique pour récupérer la mobilité de l'articulation interphalangienne proximale ont participé volontairement à cette étude. Les amplitudes articulaires ont été mesurées avant et après chaque traitement avec un goniomètre à doigt.

**Résultats.** – Chacun des deux traitements permet de récupérer les amplitudes articulaires de l'articulation interphalangienne proximale. En outre, l'efficacité de la technique surimposée est significativement supérieure à celle de la mobilisation active volontaire pour récupérer les amplitudes articulaires.

**Conclusion.** – La technique surimposée participe à une meilleure récupération des amplitudes articulaires et pourrait être intégrée aux protocoles de rééducation des entorses de l'articulation interphalangienne proximale de la main.

© 2008 Elsevier Masson SAS. All rights reserved.

**Keywords:** Neuromuscular electrical stimulation; Sprain; Hand; Proximal interphalangeal joint

**Mots clés :** Électrostimulation neuromusculaire ; Entorse ; Main ; Articulation interphalangienne proximale

---

\* Corresponding author.

E-mail address: boisgontiermatthieu@gmail.com (M. Boisgontier).

## 1. Introduction

Numerous proximal interphalangeal joint (PIP) injuries which appear to be innocuous represent actually very severe injuries to this small hinged joint and carry a high percentage of disability for the hand as well as the individual digit. Indeed, a flexion contracture of the PIP joint of one digit may significantly reduce the functional capacity of the entire hand. To prevent such physical impairment, the range of motion (ROM) must be recovered as soon as possible.

At this point, although previous studies have investigated the short-term (motor unit recruitment [6], blood flow variations [19]) and long-term (cortical excitability [8], motor recovery after stroke [15]) effects of the neuromuscular electrical stimulation (NMES), none has yet assessed, to the best of our knowledge, the possible effects of this method on recovering joints ROM [12,11].

The joint mobilization is used to limit adhesions, evacuate edema, stop the pain circle and point the healing [14,5,3]. This mobilization can either be passive, active or electrically induced. In the hand rehabilitation, the techniques of active range of motion (AROM) and of NMES are superimposed (superimposed technique [ST]: application of electrical stimulus during a voluntary muscle action) [13] in order to activate more motor units than AROM or NMES performed alone. The present study was designed to compare the efficiency of the ST to the AROM technique in improving ROM getting back. This comparison may allow us to better adapt the patient's treatment by optimizing the PIP ROM recovery after sprain.

Considering the clinical observations previously done in our rehabilitation center [17], it was hypothesized that both treatments would enhance the PIP ROM after sprain and that the ST would allow a superior ROM getting back than the treatment applying AROM on its own.

## 2. Population, materials and methods

Twenty subjects (mean age:  $36 \pm 12$  years) who were undergoing treatment within our rehabilitation center participated voluntarily in this study. They gave their informed consent to the experimental procedure as required by the Helsinki declaration (1964) and the local Ethics Committee. All subjects were cared for following a trauma with a lesion on the PIP ligaments inducing a ROM loss. The etiology of the PIP joint limitations is thus a joint contracture with passive and active ROM equal prior to treatment.

Joint ROM recovering sessions started when pain had gone, patients were cared for two times a week for 3 weeks and measurements were taken on each treatment session. Before each session, patients were randomly assigned by draw lots either to a 15 min AROM session or to a 15 min ST session. Prior to sessions, a 5 min passive mobilization session was performed. After the session of AROM or ST, 5 min tens and 5 min of massage were performed on the PIP and patients were allowed to use their hands between sessions as long as there was no pain. During sessions, the frequency of the movements was identical among both conditions and set at six finger flexions of




Fig. 1. ST sessions: patient's hand was installed on a hand rehabilitation Canadian board with electrodes onto the median nerve and the superficial flexor of fingers (SFF) muscle belly.

5 s per min with 5 s rest period after each muscle contraction. Patients used the Elpha 2000 (Danmeter A/S) stimulator for NMES [17] (Fig. 1). This device, on a 9 V tension, produced up to 60 mA contingent upon the maximum intensity tolerated by the patient. Current intensity was set prior to each session and always superior to 11 mA, that is, the mean motor threshold for muscles and the electrodes positioning of our study. The current was biphasic and asymmetric with a mean equal to zero in order to avoid any caustic damage, pulse width of 200  $\mu$ s and pulse repetition rate of 30 Hz [10,4] to involve comfortable contraction of muscles. The output was ramped over 1 s at the beginning and end of each contraction for comfort and also to attempt to prevent stretch reflexes in the antagonist muscles in response to sudden movements. This configuration is the clinical protocol applied in our rehabilitation center for many years [17].

The patient's hand was installed on a hand rehabilitation Canadian board to immobilize the metacarpalphalangeal (MCP) and wrist joints in extension [9,16] (Fig. 1) to avoid any passive ROM limitation induced by the wrist and finger extensor muscles and to make sure that the wrist and MCP joints are in the same exact position in both conditions (AROM and ST). PIP and DIP joints are free to move in maximal ROM involved by the concentric muscular contraction in AROM or ST condition.

Both sponge electrodes were placed on the skin, in a monopolar way, with an electrode interjecting an electrical stimulation onto the median nerve and the other onto the muscular belly of the superficial flexor of fingers (SFF) (Fig. 1). Exact positioning choice of the muscular electrode varied from individual to individual being based on which would get the best PIP flexion.

ROM was measured in a passive way by a physical therapist using a GEMMSOR finger goniometer, elbow at 120° flexion, wrist and MCP under extension on the hand rehabilitation Canadian board, distal interphalangeal joint in maximal extension position. The ROM measure was performed just before and just after each session.

Results obtained for all sessions were averaged for both conditions and for each subject in order to get the mean flexion ROM recovery for each treatment. The statistical test used for

Table 1  
Mean values of gains and NMES benefits in degrees.

Patients	Mean ROM gains		NMES Benefit
	AROM	ST	
1	4	12	+8
2	2	20	+18
3	4	6	+2
4	22	25	+3
5	9	12	+3
6	4	8	+4
7	4	8	+4
8	6	9	+3
9	4	6	+2
10	5	8	+3
11	3	5	+2
12	2	4	+2
13	4	6	+2
14	10	19	+9
15	3	3	+0
16	3	9	+6
17	4	10	+6
18	6	12	+6
19	13	24	+11
20	4	6	+2

the analysis was the nonparametrical test of Wilcoxon for paired samples.

### 3. Results

Analyses of the ROM measures obtained before and after each treatment showed that all subjects increased their ROM after each treatment ( $P < 0.001$ ) (Table 1).

Interestingly, ROM recovery was higher following the ST than the AROM treatment ( $P < 0.001$ ). It is to be pointed out that the mean ROM gain for all but one patient (patient 15) is better with ST than with AROM (Table 1).

### 4. Discussion

The aim of this study was to compare the effects of the AROM and ST treatments on the ROM recovery at the PIP following sprain. To address this objective, 20 patients in need of physical therapy to recover PIP ROM participated voluntarily. ROM at the PIP were measured before and after each session with a finger goniometer.

Our results showed that both treatments enhanced ROM recovery at the PIP. What is more, the ST treatment has a superior effect than AROM to enhance ROM of the PIP following ligament injury.

One of the most important parameters to take care and to discuss in this study is the NMES. At the first session of ST, although all subjects were surprised by the new sensation NMES produced, they all tolerated it. None of them dropped out once they felt stimulation. From the second session, all of them were already accustomed. The only difference remaining between the subjects was the intensity level of the stimulation they could handle. We had neither significant problem of pain and edema

related by the subjects nor any negative side effects from either treatment.

The mechanism of muscular synergies could partially explain these results. During a voluntary contraction, the agonist muscle produces the movement, while the antagonist one controls and stabilizes joints that need to remain stable [1]. This mechanism works through the myotatic reflex and the central motor patterns [13,7]. For instance, when the SFF contracts itself to bend the PIP, the intrinsics and the finger extensors contract themselves as well to control the movement and stabilize the wrist and MP joints [20,2]. In the traumatic context of edema, pain or fibrosis, this synergy enhances resistances to the movement and limits the development of maximal ROM. The NMES modifies the balance of this synergy, allowing the agonist to contract itself analytically. This analytical contraction allows an enhancement of mobility, enable to reach a better ROM which could have a cyclic effect by draining edema and therefore accentuating ROM anew.

Thus, in a future study, it would be interesting to measure the effects of the ST treatment on edema that could be an explanation of the better ROM recovery observed consecutive to this treatment. Studies on animals have shown a direct effect of electrostimulation on the edema formation after distortion [18]. Future research should also investigate the applicability of these findings to other pathologies responsible for hand or other joints ROM restrictions and for how long the ROM gains are maintained.

### References

- [1] Brand PW. Biomechanics of balance in the hand. *J Hand Ther* 1993;6:247–51.
- [2] Breteler MD, Simura KJ, Flanders M. Timing of muscle activation in a hand movement sequence. *Cereb Cortex* 2007;17:803–15.
- [3] Cannon NM. Rehabilitation approaches for distal and middle phalanx fractures of the hand. *J Hand Ther* 1993;6:285–9.
- [4] Crépon F. *Electrophysiothérapie et rééducation fonctionnelle*. 3th ed. Paris: Frison-Roche; 2002.
- [5] Chinchalkar SJ, Gan BS. Management of proximal interphalangeal joint fractures and dislocations. *J Hand Ther* 2003;16:117–28.
- [6] Feiereisen P, Duchateau J, Hainaut K. Motor unit recruitment order during voluntary and electrically induced contractions in the tibialis anterior. *Exp Brain Res* 1997;114:117–23.
- [7] Gregory CM, Bickel CS. Recruitment patterns in human skeletal muscle during electrical stimulation. *Phys Ther* 2005;85:803–15.
- [8] Khaslavskaja S, Ladouceur M, Sinkjaer T. Increase in tibialis anterior motor cortex excitability following repetitive electrical stimulation of the common peroneal nerve. *Exp Brain Res* 2002;145:309–15.
- [9] Lieber RL, Amiel D, Kaufman KR, Whitney J, Gelberman RH. Relationship between joint motion and flexor tendon force in the canine forelimb. *J Hand Surg [Am]* 1996;21:957–62.
- [10] Low J, Reed A. *Electrotherapy explained*. In: Principles and practice. Oxford: Butterworth-Heinemann; 1990.
- [11] Michlovitz SL, Harris BA, Watkins MP. Therapy interventions for improving joint range of motion: a systematic review. *J Hand Ther* 2004;17:118–31.
- [12] Michlovitz SL. Is there a role for ultrasound and electrical stimulation following injury to tendon and nerve? *J Hand Ther* 2005;18:292–6.
- [13] Paillard T, Noé F, Edeline O. Neuromuscular effects of surimposed and combined transcutaneous electrical stimulation with voluntary activity: a review. *Ann Readapt Med Phys* 2005;48:126–37.

- [14] Pettengill KM. The evolution of early mobilization of the repaired flexor tendon. *J Hand Ther* 2005;18:157–68.
- [15] Powell J, Pandyan AD, Granat M, Cameron M, Stott DJ. Electrical stimulation of wrist extensors in poststroke hemiplegia. *Stroke* 1999;30:1384–9.
- [16] Tanaka T, Amadio PC, Zhao C, Zobitz ME, An KN. Flexor digitorum profundus tendon tension during finger manipulation. *J Hand Ther* 2005;18:330–8.
- [17] Thomas D, Vaudaux G, Berthelot J. Reeduction of ligaments injuries of hand. *Cah Kinesither* 1985;114:37–53.
- [18] Thornton RM, Mendel FC, Fish DR. Effects of electrical stimulation on edema formation in different strains of rats. *Phys Ther* 1998;78:386–94.
- [19] Vanderthommen M, Depresseux JC, Dauchat L, Degueldre C, Croisier JL, Crielaard JM. Blood flow variation in human muscle during electrically stimulated exercise bouts. *Arch Phys Med Rehabil* 2002;83:936–41.
- [20] Weiss EJ, Flanders M. Muscular and postural synergies of the human hand. *J Neurophysiol* 2004;92:523–35.