A novel robotic reaching task to advance the assessment of approach-avoidance tendencies through kinematic analysis

Kayne Park^{1,2*}, Matthieu P. Boisgontier^{1,2,3*}

¹School of Rehabilitation Sciences, Faculty of Health Sciences, University of Ottawa, Canada ²Perley Health Centre of Excellence in Frailty-Informed Care, Ottawa, Canada ³Bruyère Health Research Institute, Ottawa, Canada

*Corresponding authors: apark2@uOttawa.ca (Park K); matthieu.boisgontier@uOttawa.ca (Boisgontier MP)



Abstract

The investigation of automatic approach-avoidance tendencies has traditionally relied on computer-based technologies that primarily characterise human behaviour through reaction times. However, these technologies are unable to accurately quantify other kinematic variables such as hand speed and movement direction. To address these limitations, novel robotic devices have been

developed, providing more diverse and accurate quantitative assessment of human behaviour. This technical report presents an adaptation of the approach-avoidance task on the Kinarm, a robotic platform designed to track upper limb movements as participants interact with a virtual environment. This variant of the approach-avoidance task assesses the movement of both arms in twelve directions of reach. In addition, resistive loads can be applied to investigate the role of physical effort in approach-avoidance tendencies or to support rehabilitation protocols. Data and analyses from a pilot sample (n = 5) highlights the capabilities of the Kinarm Approach-Avoidance Task (KAAT).

Keywords

Approach-Avoidance, Robotics, Reaction time, Speed, Automatic tendencies

Cite this peer-reviewed preprint:

Park K., & Boisgontier M.P. (2025). A novel robotic reaching task to advance the assessment of approach-avoidance tendencies through kinematic analysis. SportRxiv, Version 3. https://doi.org/10.51224/SRXIV.446

1. Introduction

How we move and interact with our environment can be guided by automatic tendencies (Marteau et al., 2012). For example, studies have shown that individuals approach images of physical activity faster than sedentary activities (Cheval et al., 2014; Cheval et al., 2018; Cheval et al., 2020; Farajzadeh et al., 2023a; Farajzadeh et al., 2024; Goubran et al., 2025). This bias in approaching and avoiding certain stimuli has been demonstrated in several contexts including food intake (Kemps et al., 2013; Lender et al., 2018), object ownership status (Barton et al., 2021; Roy et al., 2023), and facial expressions (Marsh et al., 2005). In addition, these approachavoidance tendencies have been associated with behaviours detrimental to health, such as physical inactivity, smoking, overeating, and alcoholism (Cheval et al., 2020; Spruyt et al., 2013; Wiers et al., 2010; Wittekind et al., 2015, 2021).

A common method for assessing an individual's approach-avoidance tendencies is the approach-avoidance task (Phaf et al., 2014; Solarz, 1960). A variant of this task has been used in studies investigating physical activity behaviour using images of physical and sedentary activities (Cheval et al., 2018; Farajzadeh et al., 2023a; Farajzadeh et al., 2024). Results consistently showed a bias: Approach responses were generated earlier for images of physical activity than for images of sedentary activity, whereas avoidance responses were earlier for sedentary activity than for physical activity, regardless of age (Farajzadeh et al., 2023a). Importantly, this variant used neutral stimuli to account for potential generic biases towards approach and avoidance movements (e.g., faster avoidance responses regardless of stimulus) (Farajzadeh et al., 2023a).

In approach-avoidance tasks, interfaces such as joysticks, mice, and keyboards are commonly used to characterise behavioural performance based on reaction times (Krieglmeyer & Deutsch, 2010; Phaf et al., 2014; Wittekind et al., 2021). However, other kinematic measures such as movement speed and acceleration may contribute to a better understanding of approach-avoidance tendencies. A good example is the study by Barton et al. (2021), who used the approach-avoidance task to examine a potential bias towards self-owned objects. They found a significant bias in reaction time for both congruent (approaching self-owned or avoiding experimenter-owned objects) and incongruent (approaching experimenter-owned or avoiding self-owned objects) mappings. However, this bias in movement duration (i.e. distance over time - speed) was only significant for the congruent mapping. Therefore, examining different measures of movement can provide a more complete characterisation of the influence that biases can have on behaviour.

In this technical report, we describe the Kinarm Approach-Avoidance Task (KAAT), a variant of the approach-avoidance task, that leverages the capabilities of robotics to better assess behaviour. The KAAT has been developed for the Kinarm Endpoint system to provide a wealth of kinematic data to characterize approach-avoidance tendencies. The stimuli used in this task can be chosen according to the objectives of a study (e.g., approach-avoidance bias for smokingrelated images, alcohol-related images, high vs. low-calorie food images, gambling-related images). Here, for the sake of illustration, we used stimuli depicting physical activities (e.g., running, swimming) and sedentary activities (e.g., lounging in a hammock, watching TV). Participants are given cues to either quickly reach towards the stimulus location ('Approach') or to reach in the opposite direction ('Avoid'). The KAAT assesses both arms and quantifies centreout reaching in 12 different directions to dissociate approach-avoidance tendencies from the bias towards specific arm motions, such as flexion versus extension (Cacioppo et al., 1993) or pushing versus pulling (Chen & Bargh, 1999; Wittekind et al., 2021). Resistive loads can be applied during reaches toward certain visual stimuli for the experimental manipulation of physical effort during reaching and for rehabilitation protocols aimed at strengthening or weakening specific approachavoidance tendencies (Farajzadeh et al., 2023b; Marteau et al., 2012).



Figure 1. (**A**) The Kinarm Approach-Avoidance Task (KAAT) on the Kinarm Endpoint laboratory. (**B**) A trial in the KAAT for the Neutral Stimulus condition with a Square Stimulus appearing in the west direction and the open circle appearing in the opposite direction (east). The dashed circles are didactic to indicate possible target locations and are not visible to the participant. The virtual hand is not visible to the participant during the task as participant's real hand is occluded by the opaque screen.

2. Kinarm Endpoint laboratory

Since the 2000s, robotic technologies have been developed to improve the quantification of human behaviour in research and clinical settings (e.g., Howard et al., 2009; Reichenbach et al., 2013; Shirota et al., 2017; Zbytniewska et al., 2019). One such technology, the Kinarm Endpoint Laboratory, is specifically designed to measure upper limb movements. This platform is a haptic robot with handles that participants can grasp and move along the horizontal plane in an augmented reality environment (Scott, 1999) (Figure 1A). A screen is positioned above the handles and provides participants with information about the position of their hands along the same horizontal plane. Participants' movements and forces are tracked by motors attached to the handles via mechanical linkages providing kinematic data. This robotic interface provides greater degrees of freedom for participants than joysticks, mice, or keyboards. Custom tasks have been developed to assess behaviour in healthy participants and to identify impairments across various conditions, such as stroke, Parkinson's disease, kidney disease, and epilepsy (Bourke et al., 2016; Gaprielian et al., 2022; Lowrey et al., 2022; Simmatis et al., 2020; Vanderlinden et al., 2022). Thus, this robotic device has proven to be an effective tool for quantifying behaviour and cognitive function.

3. Task Layout

The KAAT was developed on MATLAB (The MathWorks, Inc., version 2019b) and tested on Dexerit-E software (version 3.10, Kinarm, Kingston, Ontario). This variant of the approachavoidance task employs a 12-target centre-out reaching design (Figure 1B) to assess upper limb movements in response to visual stimuli. 24 images were used as visual stimuli (Figure 2). 12 experimental images ('Experimental Stimulus') categorized as follows: 6 images of physical activities ('Active Stimulus'; Figure 2A), 6 images of sedentary activities ('Sedentary Stimulus'; Figure 2B). 12 neutral images ('Neutral Stimulus'): 6 images made of ellipses ('Circle Stimulus'; Figure 2C) and 6 images made of rectangles ('Square Stimulus'; Figure 2D) that were designed to match the number and shape in three physical activity images (swimming, hiking, cycling) and three sedentary images (couch, hammock, reading). 2 different actions were tested ('Movement'): reaching towards a stimulus ('Approach') and reaching towards the open circle in the opposite direction ('Avoid'). Both arms were assessed in a single session. The visual stimulus could appear in 1 out of 12 possible target locations (Figure 1B). At the beginning of each block, instruction images informed the participants about the Movement-Stimulus pairing for the upcoming block (Figure 3).



Figure 2. (**A**) Images depicting physical activities used as 'Active Stimulus'. (**B**) Images depicting sedentary activities used as 'Sedentary Stimulus'. (**C**) Images used as 'Circle Stimulus'. (**D**) Images used as 'Square' Stimulus'. Each image label corresponds to the filename and each image number corresponds to the Trial Protocol used in Dexterit-E.



Figure 3. Instruction images shown to participants before (**A**) an Approach Active Stimulus block, (**B**) an Approach Sedentary Stimulus block, (**C**) an Approach Circle Stimulus block, and (**D**) an Approach Square Stimulus block.



Figure 4. Flowchart depicting the operation of the KAAT. Arrow direction indicates the next step in the task. Dashed arrows represent alternative outcomes.

4. Task operation

4.1. Sequence of events in the Kinarm Approach-Avoidance Task

The task begins with the display of an instruction image, after which the participant places the cursor of either hand in a specified target (Figure 5A). When the 'Start Target' appears, the participant moves their hand to it. Peripheral targets are then displayed, prompting the participant

to respond. If the participant does not reach a target within 3000 ms, the trial ends. If they successfully reach a target within the allotted time, the task continues as either an 'Approach Trial' or an 'Avoid Trial'. On incorrect trials (i.e., reaching a Circle Stimulus when the instruction was to approach the Square Stimulus), the text 'Incorrect' is displayed before the trial ends. The system checks whether the trial is the last in the block. If it is not, the task continues to the next trial. If it is the last trial, another check determines whether it is the final block. If the final block has been completed, the KAAT ends. Otherwise, the task progresses to the next block. This structured sequence of actions, decision points, and feedback mechanisms is illustrated in Figure 4. The following sections describe this sequence in more detail.

4.2. Starting a block

The task must be run with the 'Arm to be assessed' as 'Right'. After starting the task, participants are shown 1 out of 4 instruction images informing them of the condition (Figure 3). The block begins once the participant places either cursor (represented by a white circle of 0.5 cm radius) into the open green circle (1 cm radius) at the center of the workspace under the text 'Place either hand in the circle to start' (Figure 5A).



Figure 5. (**A**) Image taken at the start of a block. This block displays the Approach-Circle (Avoid-Square) instruction with an open green circle available for participants to move the cursor of either hand in to begin. (**B**) Start of a trial. This trial displays the blue Start target in the left side of the workspace and the participant placing their left arm into to begin the trial.

4.3. Starting a trial

At the beginning of each trial, a Start target (blue circle of 1 cm radius) appeared on the left or right side of the workspace with an equal probability (Figure 5B). After the participant placed the appropriate cursor (i.e., blue target on the left side of the workspace requires the left-hand cursor) on the Start target and held this position for 500-750 ms (randomised time), a stimulus (2.5 cm radius circle) appears in 1 of the 12 possible locations 10 cm from the Start target. Location 1 is east of the Start target, and the location for each subsequent target increases 30 degrees counterclockwise from this location (Location 4 is north, 7 is west, and 10 is south). In addition, a target (open circle of 2.5 cm radius) appears in the opposite location (i.e., if the target is in Location 1, the open circle is in Location 7). If participants do not hold the Start target for the 500-750 ms, they must re-enter the Start target and, again, hold it for 500-750 ms for the peripheral targets to appear. If participants did not reach the Start target within 3000 ms, the trial is marked as timed out, a yellow text 'Too slow' is displayed, and the trial ends.



Figure 6. Illustration of trials of an Approach Circle block. (A) Left hand correctly approaching the Circle Stimulus (white arrow not shown to participants). (B) Left hand correctly avoiding the Square Stimulus by reaching towards the open circle (white arrow not shown to participants).
 (C) Incorrect reach to the open circle resulting in an 'Incorrect!' text. (D) No reach towards any target resulting in a 'Too slow!' text.

4.4. Reaching

An approach trial is successful when the cursor is placed in any portion of the appropriate Experimental Stimulus (Figure 6A). An avoidance trial is successful when the participant's cursor moves into any portion of the open circle (Figure 6B). On an incorrect trial (e.g., cursor inside the

Square Stimulus when the instruction was to approach the Circle Stimulus), the red text 'Incorrect!' appears (Figure 6C). If the participant does not reach to any peripheral target within 3000 ms, the yellow text 'Too slow!' appears (Figure 6D). After 1100 ms of a correct trial and 1500 ms of an incorrect trial, a new Start target appears for the next trial to begin.

The KAAT has a function to apply a mechanical force to the arm when the cursor is within a certain distance of a stimulus. For example, the robot can apply a resistive load of 1 N within a 9-cm radius of the target (Figure 7).

4.5. End of a block

After 48 trials, the block ends, a new block starts, and instruction image appears.



Figure 7. Illustration of the mechanical force that can be applied to the hand (blue arrows) if the participant moves their cursor (small white circle) within a 9-cm radius from the middle of the Square Stimulus (dashed blue circle). The arrows and dashed circle are not shown to the participant.

5. Task specifications

5.1. General tab

In the General tab (Supplementary Material 1), the 'Customized instructions' in the 'Task instructions' has an explanation of the task for the participant that will display on the examiner's screen after pressing the play button. In the 'General protocol information' section, the Code is set to KAAT (Kinarm Approach Avoidance Task). 'Hand feedback behaviour' should be set to Both hands. 'Pause/unpause behaviour' should be set to Pause at next trial. 'Advanced options' should have both 'Use repeat trial flag' and 'Use random seed' checked on.

5.2. Target Table tab

In the Target Table tab (Supplementary Material 2), we assign all the relevant targets and labels used in the task.

5.2.1. Target assignment

Target 1 is a circle used for the Start target.
Target 2 is a circle used for the Avoid target.
Target 3 is a circle used to start a block.
Target 4 is a label used to display text.
Target 5 is a rectangle that displays the instruction image.
Targets 6-11 are circles that display Active image targets.
Targets 12-17 are circles that display Sedentary image targets.
Targets 18-23 are circles that display Circle image targets.
Targets 24-29 are circles that display Square image targets.

5.2.2. Target specifications

Targets 1-3 and 6-29 use Visual Radius for their circle size.

Targets 2 and 3 use Next Colour for their outline colour.

Target 4 uses Initial Colour, Next Colour, and Third Colour for label colours.

Target 5 is the instruction image target that can display 1 of 4 images found in Initial Colour (Approach-Active), Next Colour (Approach-Sedentary), Third Colour (Approach-Circle), and Fourth Colour (Approach-Square).

Targets 1, 6-29 use Initial Colour to display their images/colours.

The X column is left empty as we manually assign this value in the Task Wide Parameters tab (see below). The Visual Radius only applies to circle targets. The Target X Length and Target Y Length only apply to rectangle targets.

5.3. Trial Protocol (TP) Table tab

The Trial Protocol (TP) Table (Supplementary Material 3) assigns the targets used for the task and specifies which stimulus has a forcefield around them (Toggle Field). The first five columns each point to a single target that is consistent for all TP (e.g., all of column 1 points to target 1, all of column 2 points to target 2, etc.). The Approach Target column points to Targets 6-29 to display the different images. The CS or AS column is set to 0 to identify which TP are

Circle and Square images or 1 to identify the Active and Sedentary images. The Toggle Field column can be set to either 0 to apply no forcefield or 1 to apply a forcefield to the targets listed in the Approach Target column. In the current KAAT, the forcefield is applied to Sedentary and Square images.

5.4. Task Wide Parameters tab

The parameters set in the Task Wide Parameters tab (Supplementary Material 4) dictate the layout and flow of the task. *Avoid Distance Adjustment* scales the distance needed to move to the Avoid target (0-100%). *Distance From Midline* adjusts the distance (cm) that the Start target and peripheral targets appear relative to the midline (middle of workspace is 0 cm). *Error Delay* refers to the duration (ms) for which the 'incorrect' or 'timed out' labels are displayed. *Hold Delay* is the duration (ms) for which the cursor must remain at the Start target for a peripheral target to appear. *Maximum Force* is the strength of the forcefield (N). *Maximum Background Force* is the strength of background forces that push the hands towards the midline (N). No background load is currently used. *Ramp Duration* is the time (ms) given for forcefields to ramp up and down to the Maximum Force (half the duration is ramp up and half is ramp down). *Reach Time* is the maximum allotted time (ms) for reaching the peripheral target. *Starting Delay* is a random value from 0 to the value in Start Delay (ms) added to Hold Delay. *Success Delay* is the time (ms) the peripheral target to the centre of the peripheral or the avoid target. *Avoid Distance Adjustment* scales this value for the avoid target. *Trial Time* is the time (ms) allowed to reach the Start Target before timing out.

5.5. Block Table tab

The Block Table (Supplementary Material 5) must follow strict guidelines. First, a single block must contain the same Stimulus [only Neutral TP 13-24 (Circle/Square) or only Experimental TP 1-12 (Active/Sedentary)]. Second, the TP list must contain 12 values. Last, the value for List Reps must be 4 and Block Reps must be 1 and 'Randomized' must be checked on.

To assess all Movement-Stimulus pairings, each Stimulus must be run twice. The Movement-Stimulus pairings are assigned as follows: 1) The first instance of a Stimulus will be randomly assigned 1 of the 2 Movements (50% chance for Approach or Avoid), 2) The next block of the same Stimulus will have the opposite Movement. An example of a task sequence for 4 blocks can be as follows: Block 1 is Approach-Circle, Block 2 is Approach-Active, Block 3 is Approach-Sedentary, and Block 4 is Approach-Square.



Figure 8. Hand paths and hand speeds of an exemplar performing the KAAT. The two columns on the left display KAAT performance for Approach trials and the two columns on the right display Avoid trials. The central column illustrates the stimuli the participants were instructed to approach or avoid. Hand paths for each direction of reach are shown in the 2nd and 4th columns. Hand speed is displayed in columns 1 and 5. Data is displayed from the time period that the Approach and Avoid targets appeared (0 ms) until the end of the trial (correct target reached). Each plot presents data of the left and right arms grouped together.

6. Pilot data

Hand paths and speeds of a participant performing the KAAT are shown in Figure 8. The participant displayed relatively straight hand paths towards each direction of reach, and they displayed variability in speed from trial to trial. There was a general trend for reaches to be initiated faster for Neutral Stimulus (Circle and Square) than for Active Stimulus and Sedentary Stimulus.

Pilot data from a sample of 5 individuals (1 female, 4 males, mean age = 31.1 ± 6.7 years, age range = 28-45 years) were used to compute kinematic parameters (Table 1). Reaction Time was calculated as the difference between the time of peripheral target appearance and movement

onset, defined as the first point in time the hand speed reached 10% of the Maximum Speed for each trial. Maximum Speed was calculated as the highest hand speed from peripheral target appearance to trial end. Maximum Acceleration was calculated as the highest hand acceleration from peripheral target appearance to trial end. Speed Peaks were identified as each point in time where the hand speed was higher than both the previous and subsequent time points. Errors were identified as a trial in which the participant placed their cursor inside the incorrect stimulus (e.g., cursor in the Circle Stimulus during an Avoid Circle block) divided by the total amount of trials performed. To control for a potential generic approach or avoidance bias (Farajzadeh et al., 2023a), we subtracted the participant's mean value for the Approach Neutral Stimulus condition from the values of the Approach Active Stimulus and the values of the Approach Sedentary Stimulus condition. Similarly, the participant's mean value for the Avoid Neutral Stimulus condition was subtracted from each trial of the Avoid Active Stimulus and Avoid Sedentary Stimulus condition.

If our study were based on a larger sample of participants, results based on Reaction Time differences might suggest a bias towards approaching physical activity, as participants were faster at approaching than avoiding images depicting physical activity (110 vs. 117 ms) (Table 1). In contrast, Reaction Time differences were similar for approaching and avoiding Sedentary Stimuli, suggesting no bias towards sedentary behaviour. However, results based on speed and acceleration would challenge this latter result. Indeed, greater Maximum Speed difference, greater Speed Peak difference, and greater Maximum Acceleration difference when avoiding compared to approaching Sedentary Stimuli would suggest a bias towards avoiding Sedentary Stimuli (Table 1). These results are consistent with previous literature (Cheval et al., 2014; Cheval et al., 2018; Cheval et al., 2020; Farajzadeh et al., 2023a; Farajzadeh et al., 2024; Goubran et al., 2025).

7. Discussion

The KAAT is a bimanual multidirectional reaching task developed on the Kinarm Endpoint robot to quantify approach-avoidance tendencies towards visual stimuli. This technical report describes an overview on the images displayed in this variant of the KAAT, details of the operation to guide new users, and specifications on the backend details for users to modify the task. Pilot data and preliminary analysis of the KAAT highlight kinematic parameters that can be used to characterize approach-avoidance tendencies.

Movement	Stimulus	Mean	SD	Range	Difference Measure					
Reaction Time (ms)										
	Active	691	182	504-1297	110					
Approach	Sedentary	724	184	521-1330	143					
	Neutral	581	163	393-1170						
	Active	734	165	535-1240	117					
Avoid	Sedentary	762	190	546-1288	145					
	Neutral	617	143	447-1165						
		Maximu	m Speed (cm/s)							
	Active	54.93	8.36	38.00-75.00	2.42					
Approach	Sedentary	56.65	8.26	39.80-75.80	4.14					
	Neutral	52.51	9.87	30.20-75.80						
	Active	56.52	8.06	42.00-75.60	3.26					
Avoid	Sedentary	59.36	7.43	43.80-74.20	6.11					
	Neutral	53.26	9.37	29.40-75.60						
		Maximum A	Acceleration (cm	ו/s²)						
	Active	578	127	374-886	37					
Approach	Sedentary	629	132	381-920	88					
	Neutral	541	167	200-984						
	Active	590	140	341-935	55					
Avoid	Sedentary	647	133	410-967	111					
	Neutral	536	158	216-919						
	Reaction Time (ms) Measure Active 691 182 504-1297 110 broach Sedentary 724 184 521-1330 143 Neutral 581 163 393-1170 143 Active 734 165 535-1240 117 Active 734 165 535-1240 117 Active 734 165 535-1240 117 Active 762 190 546-1288 145 Neutral 617 143 447-1165 447-1165 Maximum Speed (cm/s) Maximum Speed (cm/s) 2.42 54.93 8.36 38.00-75.00 2.42 Active 54.93 8.36 38.07-75.80 4.14 14 14 Neutral 52.51 9.87 30.20-75.60 3.26 3.26 9.37 29.40-75.60 3.26 Sedentary 59.36 7.43 438.074.20 6.11 117 Neutral 53.26 9.37 29.40-75.60 </td									
	Active	1.09	0.30	1.00-2.20	-0.02					
Approach	Sedentary	1.16	0.40	1.00-2.60	0.05					
	Neutral	1.11	0.22	1.00-1.60						
	Active	1.06	0.21	1.00-1.80	-0.02					
Avoid	Sedentary	1.17	0.39	1.00-2.60	0.09					
	Neutral	1.08	0.31	1.00-2.60						
		E	rrors (%)							
	Active	1.04	17.89	0.00-2.10	-0.01					
Approach	Sedentary	0.63	12.25	0.00-1.05	-0.42					
	Neutral	1.05	14.49	1.04-1.05						
	Active	0.21	4.08	0.00-1.04	-0.21					
Avoid	Sedentary	1.05	14.92	0.00-3.16	0.63					
	Neutral	0.42	5.80	0.00-1.05						

Table 1. Sample	parameters	of kinematic data ((n = 5))
-----------------	------------	---------------------	---------	---

Values are shown as mean, 1 standard deviation (SD), and minimum to maximum range.

Most approach-avoidance tasks have used either a keyboard, joystick, computer mouse, or touch screen, to assess approach-avoidance tendencies towards 2 directions of movement. The KAAT quantifies approach-avoidance tendencies using 24 reach directions, increasing difficultly due to greater uncertainty in stimulus location and the motor command to be implemented, which results in longer reaction times (Burke et al., 2017; Kaufman et al., 1970; Xu et al., 2012). Further, our design included neutral stimuli, which allowed for the control of a

potential generic approach or avoidance bias (Farajzadeh et al., 2023a). Future validation work is therefore needed to examine how results in the KAAT compare with other versions of the approach-avoidance task.

Here, the KAAT was used to assess approach-avoidance tendencies towards physical activity and sedentary behaviours. However, the KAAT could be easily adapted for different goals since any image can be displayed to participants simply by changing the image files. Therefore, the KAAT can contribute to characterize health conditions such as social anxiety, depression, alcoholism, and eating disorders which has typically focused on reaction time (Heuer et al., 2007; Lender et al., 2018; Spruyt et al., 2013; Struijs et al., 2017). Further, the KAAT can refine this characterisation of human behaviour through the high sampling rate of upper limb motion - beyond what can be achieved with key presses or joystick manipulations commonly used in previous versions of the approach-avoidance task. This refined characterisation is important as some of these health conditions have been associated with sensorimotor impairments (Ben-Pazi et al., 2011; Emck et al., 2011; Northoff, 2024; Van Damme et al., 2015a; Van Damme et al., 2015b). Our pilot data highlights approach-avoidance tendencies for reaction time and movement speed. In addition, parameters such as speed peaks, which represent a change in movement direction, and acceleration, which represents movement vigor, may provide useful measures to further investigate the links between approach-avoidance tendencies, sensorimotor impairments, and health behaviours (Djamshidian et al., 2012; Kucinski et al., 2018; Mestre et al., 2013).

8. Limitations

As the aim of this technical report is to describe a novel method for assessing approachavoidance tendencies using robotics, we did not conduct hypothesis testing on our small sample size. Thus, our interpretation of the results is strictly descriptive and future work with larger samples are required to have adequate power for statistical testing.

The KAAT has been designed and tested using images of physical activity and sedentary behaviour that have been previously validated and used for an approach-avoidance task conducted on a computer-based system (Cheval et al., 2018; Farajzadeh et al., 2023a; Farajzadeh et al., 2024; Goubran et al., 2025). Whether these images effectively represent physical activity and sedentary behaviour across cultural, societal, economic, and environmental contexts remains to be examined. This potential bias of the stimuli used here to illustrate the KAAT could be mitigated by selecting other images that more accurately represent physical activity in the specific context under investigation.

15

9. Declarations

9.1. Material and code sharing

Following good research practices (Boisgontier, 2022), the material, code, and pilot data for the KAAT are freely available on the open access repository Zenodo: <u>https://doi.org/10.5281/zenodo.14879743</u> (Park & Boisgontier, 2025).

9.2. Contributions

Based on the Contributor Roles Taxonomy (CRediT) (Allen et al., 2019), individual author contributions to this work are as follows: Kayne Park: Methodology, Software, Visualization, Writing – Original Draft; Matthieu P. Boisgontier: Conceptualization, Writing – Review and Editing, Supervision, Funding Acquisition.

9.3. Funding

Kayne Park is supported by MITACS and the Banting Research Foundation (BRF 160239). Matthieu P. Boisgontier is supported by the Canada Foundation for Innovation (CFI 43661) and the Natural Sciences and Engineering Research Council of Canada (NSERC) (RGPIN-2021-03153).

9.4. Peer review and recommendation

This preprint was peer reviewed and recommended by Peer Community in (PCI) Health & Movement Sciences (Singh, 2025).

10. References

- Allen, L., O'Connell, A., & Kiermer, V. (2019). How can we ensure visibility and diversity in research contributions? How the Contributor Role Taxonomy (CRediT) is helping the shift from authorship to contributorship. *Learned Publishing*, 32(1), 71–74. https://doi.org/10.1002/leap.1210
- Barton, T., Constable, M. D., Sparks, S., & Kritikos, A. (2021). Self-bias effect: Movement initiation to self-owned property is speeded for both approach and avoidance actions. *Psychological Research*, 85(4), 1391–1406. https://doi.org/10.1007/s00426-020-01325-0

- Ben-Pazi, H., Jaworowski, S., & Shalev, R. S. (2011). Cognitive and psychiatric phenotypes of movement disorders in children: A systematic review. *Developmental Medicine and Child Neurology*, *53*(12), 1077–1084. https://doi.org/10.1111/j.1469-8749.2011.04134.x
- Boisgontier, M. P. (2022). Research integrity requires to be aware of good and questionable research practices. *European Rehabilitation Journal*, 2(1), Article 1. https://doi.org/10.52057/erj.v2i1.24
- Burke, D., Linder, S., Hirsch, J., Dey, T., Kana, D., Ringenbach, S., Schindler, D., & Alberts, J. (2017). Characterizing information processing with a mobile device: Measurement of simple and choice reaction time. *Assessment*, 24(7), 885–895. https://doi.org/10.1177/1073191116633752
- Cacioppo, J. T., Priester, J. R., & Berntson, G. G. (1993). Rudimentary determinants of attitudes:
 II. Arm flexion and extension have differential effects on attitudes. *Journal of Personality* and Social Psychology, 65(1), 5–17. https://doi.org/10.1037/0022-3514.65.1.5
- Chen, M., & Bargh, J. A. (1999). Consequences of automatic evaluation: Immediate behavioral predispositions to approach or avoid the stimulus. *Personality and Social Psychology Bulletin*, *25*(2), 215–224. https://doi.org/10.1177/0146167299025002007
- Cheval, B., Daou, M., Cabral, D. A. R., Bacelar, M. F. B., Parma, J. O., Forestier, C., Orsholits, D., Sander, D., Boisgontier, M. P., & Miller, M. W. (2020). Higher inhibitory control is required to escape the innate attraction to effort minimization. *Psychology of Sport and Exercise*, *51*, 101781. https://doi.org/10.1016/j.psychsport.2020.101781
- Cheval, B., Sarrazin, P., & Pelletier, L. (2014). Impulsive approach tendencies towards physical activity and sedentary behaviors, but not reflective intentions, prospectively predict nonexercise activity thermogenesis. *PLoS One*, *9*(12), e115238. https://doi.org/10.1371/journal.pone.0115238
- Cheval, B., Tipura, E., Burra, N., Frossard, J., Chanal, J., Orsholits, D., Radel, R., & Boisgontier, M. P. (2018). Avoiding sedentary behaviors requires more cortical resources than avoiding

physical activity: An EEG study. *Neuropsychologia*, *119*, 68–80. https://doi.org/10.1016/j.neuropsychologia.2018.07.029

- Djamshidian, A., O'Sullivan, S. S., Sanotsky, Y., Sharman, S., Matviyenko, Y., Foltynie, T.,
 Michalczuk, R., Aviles-Olmos, I., Fedoryshyn, L., Doherty, K. M., Filts, Y., Selikhova, M.,
 Bowden-Jones, H., Joyce, E., Lees, A. J., & Averbeck, B. B. (2012). Decision making,
 impulsivity, and addictions: Do Parkinson's disease patients jump to conclusions? *Movement Disorders*, *27*(9), 1137–1145. https://doi.org/10.1002/mds.25105
- Emck, C., Bosscher, R. J., Van Wieringen, P. C., Doreleijers, T., & Beek, P. J. (2011). Gross motor performance and physical fitness in children with psychiatric disorders.
 Developmental Medicine & *Child Neurology*, 53(2), 150–155. https://doi.org/10.1111/j.1469-8749.2010.03806.x
- Farajzadeh, A., Goubran, M., Beehler, A., Cherkaoui, N., Morrison, P., de Chanaleilles, M., Maltagliati, S., Cheval, B., Miller, M. W., Sheehy, L., Bilodeau, M., Orsholits, D., & Boisgontier, M. P. (2023). Automatic approach-avoidance tendency toward physical activity, sedentary, and neutral stimuli as a function of age, explicit affective attitude, and intention to be active. *Peer Community Journal*, *3*, 246. https://doi.org/10.24072/pcjournal.246
- Farajzadeh, A., Goubran, M., Fessler, L., Cheval, B., Allen, Z. V., & Boisgontier, M. P. (2023). Training older adults to inhibit the automatic attraction to sedentary stimuli: A cognitivebias-modification protocol. *European Rehabilitation Journal*, 3(1), 1-9. https://doi.org/10.52057/erj.v3i1.32
- Farajzadeh, A., Jabouille, F., Benoit, N., Bezeau, O., Bourgie, T., Gerro, B., Ouimet, J., & Boisgontier, M. P. (2024). Apathy, intentions, explicit attitudes, and approach-avoidance tendencies in physical activity behavior. *Communications in Kinesiology*, 1(6). https://doi.org/10.51224/cik.2024.68

- Gaprielian, P., Scott, S. H., & Levy, R. (2022). Reverse visually guided reaching in patients with
 Parkinson's disease. *Parkinson's Disease*, 2022(1), 8132923.
 https://doi.org/10.1155/2022/8132923
- Goubran, M., Zammar, C., Tellez Alvarez, S., Héran, E., Proulx., S, Bilodeau, M., & Boisgontier,
 M. P. (2025). Approach-avoidance tendencies influence the relationship between fear of
 movement and physical activity in osteoarthritis. *MedRxiv*, 2025.
 https://doi.org/10.1101/2025.01.23.25321044
- Heuer, K., Rinck, M., & Becker, E. S. (2007). Avoidance of emotional facial expressions in social anxiety: The Approach–Avoidance Task. *Behaviour Research and Therapy*, 45(12), 2990–3001. https://doi.org/10.1016/j.brat.2007.08.010
- Howard, I. S., Ingram, J. N., & Wolpert, D. M. (2009). A modular planar robotic manipulandum with end-point torque control. *Journal of Neuroscience Methods*, *181*(2), 199–211. https://doi.org/10.1016/j.jneumeth.2009.05.005
- Kaufman, H., Lamb, J. C., & Walter, J. R. (1970). Prediction of choice reaction time from information of individual stimuli. *Perception & Psychophysics*, 7(5), 263–266. https://doi.org/10.3758/BF03210161
- Kemps, E., Tiggemann, M., Martin, R., & Elliott, M. (2013). Implicit approach–avoidance associations for craved food cues. *Journal of Experimental Psychology: Applied*, 19(1), 30–38. https://doi.org/10.1037/a0031626
- Krieglmeyer, R., & Deutsch, R. (2010). Comparing measures of approach–avoidance behaviour:
 The manikin task vs. two versions of the joystick task. *Cognition and Emotion*, 24(5), 810–828. https://doi.org/10.1080/02699930903047298
- Kucinski, A., Lustig, C., & Sarter, M. (2018). Addiction vulnerability trait impacts complex movement control: Evidence from sign-trackers. *Behavioural Brain Research*, 350, 139– 148. https://doi.org/10.1016/j.bbr.2018.04.045

- Lender, A., Meule, A., Rinck, M., Brockmeyer, T., & Blechert, J. (2018). Measurement of foodrelated approach-avoidance biases: Larger biases when food stimuli are task relevant. *Appetite*, *125*, 42–47. https://doi.org/10.1016/j.appet.2018.01.032
- Lowrey, C. R., Dukelow, S. P., Bagg, S. D., Ritsma, B., & Scott, S. H. (2022). Impairments in cognitive control using a reverse visually guided reaching task following stroke. *Neurorehabilitation and Neural Repair*, 36(7), 449–460. https://doi.org/10.1177/15459683221100510
- Marsh, A. A., Ambady, N., & Kleck, R. E. (2005). The effects of fear and anger facial expressions on approach- and avoidance-related behaviors. *Emotion*, 5(1), 119–124. https://doi.org/10.1037/1528-3542.5.1.119
- Marteau, T. M., Hollands, G. J., & Fletcher, P. C. (2012). Changing human behavior to prevent disease: The importance of targeting automatic processes. *Science*, 337(6101), 1492– 1495. https://doi.org/10.1126/science.1226918
- Mestre, T. A., Strafella, A. P., Thomsen, T., Voon, V., & Miyasaki, J. (2013). Diagnosis and treatment of impulse control disorders in patients with movement disorders. *Therapeutic Advances in Neurological Disorders*, 6(3), 175–188. https://doi.org/10.1177/1756285613476127
- Northoff, G. (2024). Beyond mood depression as a speed disorder: Biomarkers for abnormal slowness. *Journal of Psychiatry and Neuroscience*, *49*(5), E357–E366. https://doi.org/10.1503/jpn.240099
- Park, K., & Boisgontier, M. P. (2025). A novel robotic reaching task to advance the assessment of approach-avoidance tendencies through kinematic analysis: scripts and data. *Zenodo, Version 1.0.* <u>https://doi.org/10.5281/zenodo.14879743</u>
- Phaf, R. H., Mohr, S. E., Rotteveel, M., & Wicherts, J. M. (2014). Approach, avoidance, and affect:
 A meta-analysis of approach-avoidance tendencies in manual reaction time tasks. *Frontiers in Psychology*, 5. https://doi.org/10.3389/fpsyg.2014.00378

- Reichenbach, A., Costello, A., Zatka-Haas, P., & Diedrichsen, J. (2013). Mechanisms of responsibility assignment during redundant reaching movements. *Journal of Neurophysiology*, *109*(8), 2021–2028. https://doi.org/10.1152/jn.01052.2012
- Roy, N., Karnick, H., & Verma, A. (2023). Towards the self and away from the others: Evidence for self-prioritization observed in an approach avoidance task. *Frontiers in Psychology*, *14*, 1041157. https://doi.org/10.3389/fpsyg.2023.1041157
- Scott, S. H. (1999). Apparatus for measuring and perturbing shoulder and elbow joint positions and torques during reaching. *Journal of Neuroscience Methods*, *89*(2), 119–127. https://doi.org/10.1016/S0165-0270(99)00053-9
- Shirota, C., van Asseldonk, E., Matjačić, Z., Vallery, H., Barralon, P., Maggioni, S., Buurke, J. H.,
 & Veneman, J. F. (2017). Robot-supported assessment of balance in standing and walking. *Journal of Neuroengineering and Rehabilitation*, 14(1), 80.
 https://doi.org/10.1186/s12984-017-0273-7
- Simmatis, L. E., Jin, A. Y., Taylor, S. W., Bisson, E. J., Scott, S. H., & Baharnoori, M. (2020). The feasibility of assessing cognitive and motor function in multiple sclerosis patients using robotics. *Multiple Sclerosis Journal - Experimental, Translational and Clinical*, 6(4). https://doi.org/10.1177/2055217320964940
- Singh, T. (2025) A novel Kinarm-based behavioral paradigm to probe reaching and avoidance actions. *Peer Community in Health and Movement Sciences*, 100187. 10.24072/pci.healthmovsci.100187
- Solarz, A. K. (1960). Latency of instrumental responses as a function of compatibility with the meaning of eliciting verbal signs. *Journal of Experimental Psychology*, *59*(4), 239–245. https://doi.org/10.1037/h0047274
- Spruyt, A., De Houwer, J., Tibboel, H., Verschuere, B., Crombez, G., Verbanck, P., Hanak, C., Brevers, D., & Noël, X. (2013). On the predictive validity of automatically activated

approach/avoidance tendencies in abstaining alcohol-dependent patients. *Drug and Alcohol Dependence*, *127*(1), 81–86. https://doi.org/10.1016/j.drugalcdep.2012.06.019

- Struijs, S. Y., Lamers, F., Vroling, M. S., Roelofs, K., Spinhoven, P., & Penninx, B. W. J. H. (2017).
 Approach and avoidance tendencies in depression and anxiety disorders. *Psychiatry Research*, 256, 475–481. https://doi.org/10.1016/j.psychres.2017.07.010
- Van Damme, T., Fransen, E., Simons, J., van West, D., & Sabbe, B. (2015a). Motor impairment among different psychiatric disorders: Can patterns be identified? *Human Movement Science*, 44, 317–326. https://doi.org/10.1016/j.humov.2015.10.006
- Van Damme, T., Simons, J., Sabbe, B., & van West, D. (2015b). Motor abilities of children and adolescents with a psychiatric condition: A systematic literature review. World Journal of Psychiatry, 5(3), 315–329. https://doi.org/10.5498/wjp.v5.i3.315
- Vanderlinden, J. A., Semrau, J. S., Silver, S. A., Holden, R. M., Scott, S. H., & Boyd, J. G. (2022). Acute kidney injury is associated with subtle but quantifiable neurocognitive impairments. *Nephrology Dialysis Transplantation*, 37(2), 285–297. https://doi.org/10.1093/ndt/gfab161
- Wiers, R. W., Rinck, M., Kordts, R., Houben, K., & Strack, F. (2010). Retraining automatic actiontendencies to approach alcohol in hazardous drinkers. *Addiction*, 105(2), 279–287. https://doi.org/10.1111/j.1360-0443.2009.02775.x
- Wittekind, C. E., Blechert, J., Schiebel, T., Lender, A., Kahveci, S., & Kühn, S. (2021). Comparison of different response devices to assess behavioral tendencies towards chocolate in the approach-avoidance task. *Appetite*, 165, 105294. https://doi.org/10.1016/j.appet.2021.105294
- Wittekind, C. E., Feist, A., Schneider, B. C., Moritz, S., & Fritzsche, A. (2015). The approach-avoidance task as an online intervention in cigarette smoking: A pilot study. *Journal of Behavior Therapy and Experimental Psychiatry*, 46, 115–120. https://doi.org/10.1016/j.jbtep.2014.08.006

22

- Xu, Y., Terekhov, A., Latash, M., & Zatsiorsky, V. (2012). Forces and moments generated by the human arm: Variability and control. *Experimental Brain Research*, 223(2), 159–175. https://doi.org/10.1007/s00221-012-3235-0
- Zbytniewska, M., Rinderknecht, M. D., Lambercy, O., Barnobi, M., Raats, J., Lamers, I., Feys, P., Liepert, J., & Gassert, R. (2019). Design and characterization of a robotic device for the assessment of hand proprioceptive, motor, and sensorimotor impairments. *IEEE International Conference on Rehabilitation Robotics*, 2019, 441–446. https://doi.org/10.1109/ICORR.2019.8779507

SUPPLEMENTARY MATERIAL

A novel robotic reaching task to advance the assessment of approachavoidance tendencies through kinematic analysis

Kayne Park, Matthieu P. Boisgontier

Supplementary Material 1. Screenshot of the General tab in the sandwichOrder.dtp file

Supplementary Material 2. Screenshot of the Target Table tab displaying columns used to set target properties.

Supplementary Material 3. Screenshot of the Trial Protocol (TP) Table identifying the targets to use for specific columns.

Supplementary Material 4. Screenshot of the Task Wide Params tab.

Supplementary Material 5. Screenshot of the Block Table highlighting the TP List, List Reps, Randomized, and Block Reps.

Supplementary Material 1. Screenshot of the General tab in the sandwichOrder.dtp file

Edit Task Protocol (sandwichOrder)

General Target Table Load Table TP Table Task Wide Params Block	k Table Analog Inputs Input Files
Task instructions	
Default instructions	Customized instructions This task examines how quickly you can respond to visual input. At the beginning of each block of trials, an instruction image will appear highlighting which images to approach and which to avoid. Once this association is learned, place either cursor into the open green circle to begin. Video
General protocol information Code: KAAT Description: Kinarm Approach Avoid Task	
Hand feedback behaviour Gaz Selected behaviour: Both hands Selected behaviour Feedback will be shown for both hands.	e feedback behaviour lected behaviour: Off Gaze_Feedback block not found in model.
Pause/unpause behaviour Selected behaviour: Pause at next trial Pause at the beginning of the next trial. Pressing the Run/Pause but	ton again continues the next trial.
Advanced options Use repeat trial flag Use random seed	Data rate 1000 Hz Samples per second saved

Supplementary Material 2. Screenshot of the Target Table tab displaying columns used to set

target properties.

jet lable												
X Position (m) of th:	e target i	relative to local	(0,0)								
	x	Y	Visual Radius	Target X Length	Target Y Length	Initial Colour	Next Colour	Label Text	Label Height	Stroke Width	Third Colour	Fourth Colour
arget 1	0.000	-39.000	1.000	0.000	0.000				0.000	0.000		
arget 2	0.000	-39.000	2.500	0.000	0.000				0.000	0.200		
arget 3	0.000	-39.000	1.000	0.000	0.000				0.000	0.200		
arget 4	0.000	-37.000	0.000	0.000	0.000				2.000	0.000		
arget 5	0.000	-25.000	0.000	32.000	16.000	A. S. M. G. S.	No. 12 The State of State		0.000	0.000	·····	
arget 6	0.000	-39.000	2.500	0.000	0.000				0.000	0.000		
arget 7	0.000	-39.000	2.500	0.000	0.000	13-5-5			0.000	0.000		
arget 8	0.000	-39.000	2.500	0.000	0.000				0.000	0.000		
arget 9	0.000	-39.000	2.500	0.000	0.000				0.000	0.000		
arget 10	0.000	-39.000	2.500	0.000	0.000	1-1_			0.000	0.000		
arget 11	0.000	-39.000	2.500	0.000	0.000	- Stas			0.000	0.000		
arget 12	0.000	-39.000	2.500	0.000	0.000	ليحصب يحمكم			0.000	0.000		
arget 13	0.000	-39.000	2.500	0.000	0.000	Pick- art?			0.000	0.000		
arget 14	0.000	-39.000	2.500	0.000	0.000	THE - E			0.000	0.000		
arget 15	0.000	-39.000	2.500	0.000	0.000	Clerit -			0.000	0.000		
arget 16	0.000	-39.000	2.500	0.000	0.000	125mm			0.000	0.000		
arget 17	0.000	-39.000	2.500	0.000	0.000	Cherry Fr			0.000	0.000		
	0.000	20.000	2 500	0.000	0.000	(0.000	0.000		

Target coordinates are (x,y) positions defined relative to the midpoint between the robot origins.

Supplementary Material 3. Screenshot of the Trial Protocol (TP) Table identifying the targets to use for specific columns.

ral	Target Table	Load Tabl	e TP Table	Task Wide Parar	ns Block Tab	le Analog Ing	outs Input Files		
Pro	otocol (TP) Tak	ble							
art Ta	arget: Pick the	target for	the starting tar	get					
				1					
	Star	rt Target	Avoid Target	Begin target	Label target	Info Target	Approach Target	CS or AS	Toggle Field
P 1		1	2	3	4	5	6	1	0
TP 2		1	2	3	4	5	7	1	0
TP 3		1	2	3	4	5	8	1	0
TP 4		1	2	3	4	5	9	1	0
TP 5		1	2	3	4	5	10	1	0
TP 6		1	2	3	4	5	11	1	0
TP 7		1	2	3	4	5	12	1	1
TP 8		1	2	3	4	5	13	1	1
TP 9		1	2	3	4	5	14	1	1
TP 1	0	1	2	3	4	5	15	1	1
TP 1	1	1	2	3	4	5	16	1	1
TP 12	2	1	2	3	4	5	17	1	1
TP 1	3	1	2	3	4	5	18	0	0
TP 14	4	1	2	3	4	5	19	0	0
TP 1	5	1	2	3	4	5	20	0	0
TP 1	6	1	2	3	4	5	21	0	0
TP 1	7	1	2	3	4	5	22	0	0
TP 1	8	1	2	3	4	5	23	0	0
TP 19	9	1	2	3	4	5	24	0	1
TP 2	0	1	2	3	4	5	25	0	1
TP 2	1	1	2	3	4	5	26	0	1
TP 2	2	1	2	3	4	5	27	0	1
TP 2	3	1	2	3	4	5	28	0	1
TP 24	4	1	2	3	4	5	29	0	1
TP 2	5	0	0	0	0	0	0	0	0
TP 2	6	0	0	0	0	0	0	0	0
TP 2	7	0	0	0	0	0	0	0	0
TP 2	8	0	0	0	0	0	0	0	0

Supplementary Material 4. Screenshot of the Task Wide Params tab.

Edit Task Protocol (sandwichOrder)

General	Target Table	Load Table	TP Table	Task Wide Para	ams Block Table	Analog Inputs	Input Files
- Task W	ide Parameters	s					
	Avoid Distar	nce Adjustme	ent 100.0	00	Modify the distan	ce from centre fo	or the avoid target (%) (100% means distance fr
	Distance Fro	om Midline	17.50)	Adjust the distand	e for left and rig	yht start point from midline (cm)
	Error Delay		750.0	00	Time after an erro	or trial (ms)	
	Hold Delay		500.0	00	Time to hold star	t target to begin t	trial (ms)
	Maximum F	orce	1.00		Maximum amour	nt of force allowe	ed on position controller (robot limits to 30N wł
	Maximum F	orce Backgro	und 0.00		Maximum force a	llowed for backg	ground load (N)
	Ramp Durat	ion	100.0	00	Time for forces to	ramp up or dow	vn (ms)
	Reach Time		3000	.00	Maximum amour	nt of time to reac	ch for peripheral targets (ms)
	Starting Dela	ау	250.0	00	Randomly genera	te a value from 0	0 to Starting Delay to add to hold delay (ms)
	Success Dela	ау	300.0	00	Time after a succ	essful trial (ms)	
	Target Dista	nce	10.00)	Distance from cer	ntre target to cent	ntre of peripheral targets (cm)
	Trial Time		3000	.00	Maximum amour	nt of time to start	t trial (ms)

Supplementary Material 5. Screenshot of the Block Table highlighting the TP List, List Reps, Randomized, and Block Reps.

iai iaigei	Table Load Table TP Table Task Wide Params Blo	ck lable Analog Inp	uts Input Files		
ck Table —					
	TP List	List Reps	Catch TP List	Randomized	Block Reps
Block 1	13-24	4			1
Block 2	1-12	4			1
Block 3	1-12	4			1
Block 4	13-24	4		\checkmark	1
Block 5		0			0
Block 6		0			0
Block 7		0			0
		0			0
Block 8		0			0
Block 8 Block 9		• •			
Block 8 Block 9 Block 10		0			0